

Hale COLLAGE problem set 1 (Due Wed. Feb 15)

- The first lectures used simple problems, formulated in two dimensions, to illustrate the role of magnetic reconnection in releasing stored energy. Here we explore such model for a compact flare — no flux rope or eruption. The model involves two identical magnetic bipoles, whose field is shown in fig. 1a. Any magnetic field in the (y, z) plane, can be written using a flux function $A(y, z)$

$$\mathbf{B}(y, z) = \nabla A \times \hat{\mathbf{x}} = \frac{\partial A}{\partial z} \hat{\mathbf{y}} - \frac{\partial A}{\partial y} \hat{\mathbf{z}} . \quad (1)$$

Contours of the function show field lines, and its value at any point ($z \geq 0$) gives the flux (per ignorable length) ψ passing between that point and the origin $(y, z) = (0, 0)$. (If you have no experience working with flux functions, you might try convincing yourself of these important properties using vector calculus.)




Figure 1: The field above a pair of line-bipoles $P1-N1$ (left) and $P2-N2$ (right). (a) Shows the potential field defined by eq. (2). (b) Shows the field with a single island, given by eq. (3), used as a simple model of a current sheet. The flux linking $P1$ to $N2$, denoted ψ_{12} is the same in both examples.




Figure 2: The field when the inner separation, $2a$, is decreased (left) and increased (right). The null point has been deformed into a current sheet, shown with a dark curve. The flux ψ_{12} connected $P1$ to $N2$ is the same in each case as in the potential field, fig. 1a.

The pair of bipoles consists of four sources, two positive and two negative, located on the photosphere ($z = 0$) and arranged symmetrically about the z axis ($y = 0$) at points

$y = \pm a$ and $y = \pm b$ (see fig. 1a). The potential field, $\nabla \times \mathbf{B} = 0$, above this distribution is generated by the flux function

$$A^{(p)}(y, z) = \lambda \tan^{-1} \left(\frac{y+b}{z} \right) - \lambda \tan^{-1} \left(\frac{y+a}{z} \right) + \lambda \tan^{-1} \left(\frac{y-a}{z} \right) - \lambda \tan^{-1} \left(\frac{y-b}{z} \right), \quad (2)$$

for $z > 0$, where the magnetic flux (per ignorable length) of each sources in $\pi\lambda$. (You can see from the definition that $A^{(p)} = \pi\lambda$ at points on the photosphere $-b < y < -a$ and $a < y < b$.)

- The potential field has a single X-point at $(y, z) = (0, z_x)$. Find the value of z_x explicitly in terms of a , b , and λ . Evaluate the flux function at that point to obtain the amount of flux, $\psi_{12}^{(p)}$, connecting $P1$ to $N2$ in a potential field. Verify that this expression has the correct values in limits $a/b \rightarrow 1$ and $a/b \rightarrow 0$.
- Say we begin with $a = b/2$, and then change a by a small amount $\Delta a \ll a$. Compute the change $\Delta\psi_{12}^{(p)}$ to lowest order in $\Delta a/a \ll 1$.
- In an ideal plasma, where $E' = 0$, the flux connecting $P1$ to $N2$, ψ_{12} , will not change. If we change a , as in part *b*, the field will develop a current sheet in place of the null point, as shown in fig. 2. It is possible to approximate the sheet with a set of wires whose islands link together to form a chain. The simplest such model has a *single* wire, located at $z = h$, and carrying current I_{cs} . This creates a single island, as shown in fig. 1b. The field is given by a potential

$$A(y, z) = A^{(p)}(y, z) + \frac{I_{cs}}{c} \ln \left[\frac{y^2 + (z+h)^2}{y^2 + (z-h)^2} \right], \quad (3)$$

where $A^{(0)}(y, z)$ is given by eq. (2). (Note that the added term vanishes along $z = 0$, so the wire does not affect the vertical photospheric field. This is because the added terms consists of a wire, at $z = h$, and an oppsing *image current* at $z = -h$.)

The field generated by eq. (3), contains two null points in place of the single null in $A^{(0)}(y, z)$. To model a current sheet these *must* occur at the same value of A , as in fig. 1b. This value gives the flux ψ_{12} in the presence of the current sheet. The separation between the two nulls approximates the extent of the actual current sheet, $2L$, as in fig. 2.

For which sign of I_{cs} will these null points both fall on the z axis as in fig. 1b? Explain your answer in words. Use this sign and find the null positions in the limit of small current. In this limit you may take $h = z_x$, from part a., and expand expressions in powers of $z - z_x$, to obtain a location to leading order in $I_{cs}/c\lambda \ll 1$. Find both nulls and verify that they each have the same value of A (justifying our assignment of $h = z_x$).

- Use the results of c. to compute the flux difference from a potential field in the case $b = 2a$ to leading order in I_{cs} . This should take a form

$$\Delta\psi_{12} = \psi_{12} - \psi_{12}^{(p)} \propto I_{cs} \ln(\alpha |I_{cs}|), \quad (4)$$

where you need to find the constant of proportionality *and* the value of α . Next find the distance between X-points, as an approximation to the full length of the current sheet, $2L$. Find this to leading order in current, explicitly in terms of I_{cs} , λ and a .

- e. The energy (per ignorable length) released by complete reconnection of a current sheet ($\Delta\psi_{12} \rightarrow 0$) can be found from the electromagnetic work integral

$$\Delta\mathcal{E}_M = \frac{1}{c} \int_0^{\Delta\psi_{12}} I_{cs}(\Delta\psi_{12}) d(\Delta\psi_{12}) . \quad (5)$$

Use the approximate expression from part d. to perform the integral explicitly and obtain an explicit expression in terms of I_{cs} . (This might require an integration by parts.)

- f. We may use this simple model to obtain the magnetic field strength B_i just outside the actual current sheet. Ampère's law,

$$\frac{4\pi|I_{cs}|}{c} = \oint \mathbf{B} \cdot d\mathbf{l} \simeq 4L B_i , \quad (6)$$

can be used, in conjunction with the length and current from e., to obtain an explicit expression for B_i in terms of I_{cs} , λ and a , for the case $b = 2a$. Assuming some uniform mass density ρ_0 write down the value of the Alfvén-transit time

$$\tau_A = \frac{2L}{v_A} , \quad (7)$$

for the current sheet.

- g. We now use the results above to obtain numerical values for a simplified model of a compact flare. Consider a case where $a = 3 \times 10^9$ cm, and $b = 2a = 6 \times 10^9$ cm. We will assign a finite extent, $L_x = 10^{10}$ cm in the previously ignorable direction, but continue to use the two-dimensional expressions obtained above. Assign the parameter λ so that every source has a total flux of 10^{22} Mx. The current sheet builds up as the inner sources each move by $\Delta a = 10^9$ cm under *ideal* conditions ($E' = 0$). Assume the motion is in the direction which produces a *vertical* current sheet. Use the expressions derived for $\Delta a \ll a$ to find the length $2L$ of the resulting current sheet, the height of its center, h , the current it carries $|I_{cs}|$ (express this in Amps), and the energy available for release by magnetic reconnection. Finding the current will require you to solve a transcendental equation. You may do this approximately in whichever manner you prefer.
- h. Following the storage phase (part g.) there is sudden and complete reconnection of the current sheet, restoring a potential field ($\Delta\psi_{12} \rightarrow 0$). This produces flare ribbons in a region whose photospheric, vertical field strength is $B_{z,0} = 300$ G (use this instead of the pathological values you would obtain using eq. (3) at $z = 0$). Each flare ribbon moves horizontally with a mean velocity $v_{\text{rib}} = 3$ km/s. What is the time, τ_{rx} , required for complete reconnection? Use this to compute the average power released in the flare, the mean reconnection electric field (expressed in V/m), and the energy flux incident on each ribbon (erg/cm²/s). Finally assume a mass density $\rho_0 = 10^{-15}$ g/cm³ at the current sheet and compute the Alfvén Mach number of the reconnection.